

# PROCEEDINGS

OF THE

# BIOLOGICAL SOCIETY OF WASHINGTON

### MUTANDA ORNITHOLOGICA.

III.

#### BY HARRY C. OBERHOLSER.

The present paper is the third\* in the writer's series of articles concerning nomenclatorial changes in the names of birds. This treats of some alterations in specific terms in the families Haematopodidae, Scolopacidae, Loriidae, Treronidae, and Columbidae.

### FAMILY HAEMATOPODIDAE.

# Haematopus quoyi Brabourne and Chubb.

Messrs. Brabourne and Chubb have recently given the name Haematopus quoyi† to the bird from the Falkland Islands previously described by Sharpe‡ as Haematopus ater. There is, however, an earlier and certainly pertinent name in Hoematopus townsendi Audubon,§ based on a specimen supposedly from the mouth of the Columbia River, but which doubtless came, as did others of Audubon's specimens, from South America. In view of this, we designate the Falkland Islands as its type locality. The name for this species should, therefore, now become Haematopus townsendi Audubon.

#### FAMILY SCOLOPACIDAE.

#### Totanus fuscus (Linnaeus).

The spotted redshank of Europe is now called *Totanus fuscus*, from *Scolopax fusca* Linnaeus. || This name, however, is preoccupied by *Scolopax* 

15-Proc. Biol. Soc. Wash., Vol. 31, 1918.

(47)

of Mut on vi

<sup>\*</sup>For the two previous papers, cf. Proc. Biol. Soc. Wash., XXX, March 31, 1917, pp. 75-76; and Proc. Biol. Soc. Wash., XXX, July 27, 1917, pp. 125-126.

<sup>†</sup> Birds South Amer., I, December, 1912, p. 37.

Cat. Birds Brit, Mus., XXIV, 1896, p. 121.

Birds Amer., folio ed., IV, 1838, pl. 427, fig. 2 ("mouth of the Columbia River, Oregon.").

<sup>||</sup> Syst. Nat., ed. 12, I, 1766, p. 243 (Europe).

fusca Linnaeus,\* which is apparently the same as Guara alba (Linnaeus). The earliest tenable name for Totanus fuscus of authors is apparently Scolopax maculata Tunstall,† which the author's reference to Pennant's British Zoology saves from being a nomen nudum. The species should, therefore, now stand as Totanus maculatus (Tunstall).

eryphropios

# FAMILY LORIIDAE.

### Eos riciniata (Bechstein).

The name of this lory in general use; is antedated by at least three others. Dr. R. B. Sharpes has used for it *Eos variegata* (Gmelin); but the earliest of its specific designations is *Psittacus guenbyensis* Scopoli, which, though from an impossible locality, is of apparently undoubted application. The species should therefore stand as *Eos guenbyensis* (Scopoli).

#### FAMILY TRERONIDAE.

### Leucotreron gularis (Quoy and Gaimard).

The Columba gularis of Quoy and Gaimard,\*\* which is the present Leucotreron gularis, from Celebes, is preoccupied by Columba gularis Wagler,†† which is a synonym of Coturnicoenas hottentotta (Temminck), a fictitious bird of Levaillant's. As Leucotreron gularis has no other name, it may be called Leucotreron epia, nom. nov.

#### FAMILY COLUMBIDAE.

# Gymnopelia erythrothorax (Meyen).

The name now employed for this South American species, Columba erythrothorax Meyen, ‡‡ is untenable because of Columba erythrotorax [sic] Temminck, §§ which is probably the same as Aplopelia larvata Temminck. The next available name is Columba (Chamæpelia) cecilioe Lesson || || ||, and

<sup>\*</sup> Syst. Nat., ed. 10, I, 1758, p. 145 (South Carolina).

<sup>† [</sup>Scolopax] Maculata Tunstall, Ornith. Brit., 1771, p. 3 (Great Britain).

<sup>†</sup> Ps[ittacus]. riciniatus Bechstein, Latham's Allgem. Übers. Vögel, VII, 1811, p. 69, pl. 4 (Molucca Islands).

<sup>§</sup> Hand-List Gen. and Spec. Birds, II, 1900, p. 2.

<sup>| [</sup>Psittacus] variegatus Gmelin, Syst. Nat., I, pt. 1, 1788, p. 319 (India).

T Psittacus guenbyensis Scopoli, Del. Flor. et Faun. Insubr., II, 1786, p. 87 ("China"; locality wrong; doubtless came from Molucca Islands; therefore we designate Gilolo Island as the type locality).

<sup>\*\*</sup> Voy. Astrolabe, Zool., I, 1830, p. 247, pl. 29 (Manado, Celebes).

<sup>++</sup> Syst. Ayium, 1827, Columba, sp. 90, p. 261 (Namaqua Land, South Africa).

tt Nov. Act. Acad. Leopold.-Carol., XVI, Suppl., 1834, p. 98, t. XVI (province of Arequipa, Peru).

<sup>Mist. Nat. Gén. Pigeons, I, fam. trois., 1808-1811, p. 15, pl. VII (Surinam).</sup> 

<sup>||||</sup> Echo du Monde Savant, XII, 1845, p. 8 (Peru). This was later remained Columba (Chamœpelia) anais by Lesson, in Compl. Oeuv. Buffon, XX, 1847, p. 210 (Peru).